arXiv:1612.03084v3 [math.GT] 20 Dec 2016

Is a typical bi-Perron algebraic unit a pseudo-Anosov dilatation?

HYUNGRYUL BAIK, AHMAD RAFIQI, AND CHENXI WU

Abstract. In this note, we deduce a partial answer to the question in the title. In particular, we show that asymptotically almost all bi-Perron algebraic unit whose characteristic polynomial has degree at most $2n$ do not correspond to dilatations of pseudo-Anosov maps on a closed orientable surface of genus n for $n \geq 10$. As an application of the argument, we also obtain a statement on the number of closed geodesics of the same length in the moduli space of area one abelian differentials for low genus cases.

Thurston's celebrated theorem [19] classified self-homeomorphisms of compact orientable surfaces up to isotopy. As long as a homeomorphism does not admit a finite power which is isotopic to the identity and does not admit a finite union of disjoint simple closed curves which are preserved up to isotopy, it is isotopic to a so-called *pseudo-Anosov map*. Associated to a pseudo-Anosov map is a positive number called its *dilatation*. The logarithm of the dilatation of a pseudo-Anosov map is the topological entropy of the map. See [21] for details.

We call a number λ to be a *pseudo-Anosov dilatation* if it is the dilatation constant for a pseudo-Ansov map on some compact orientable surface. We call an algebraic integer λ *bi-Perron* if all the Galois conjugates of λ are contained in an annulus of outer radius λ itself and inner radius $1/\lambda$. Fried [7] showed that every pseudo-Anosov dilatation is a bi-Perron algebraic unit, and he conjectured that this is also a sufficient condition: every bi-Perron algebraic unit is realized as the dilatation constant of some pseudo-Anosov map.

This turned out to be a difficult question and there are many related works. Thurston [20] gave an answer to an $\mathrm{Out}(F_n)$-version of this question. In [20], Thurston also gave a few examples of constructions of pseudo-Anosov maps for given bi-Perron algebraic units (here the numbers are given as the slopes of some post-critically finite piecewise-linear self-homeomorphisms of the unit interval). These examples motivated the authors' previous work to construct pseudo-Anosov maps in [2] for a bi-Perron algebraic unit given as the leading eigenvalue of a Perron-Frobenius matrix satisfying some additional properties. Both constructions in [20] and [2] are similar to the one given in [3]. See also, for instance, [1], [15], [17], [14], [11], [12], [13], [18] for various constructions of pseudo-Anosov maps.

In this short note, we formulate an easier version of Fried's question, and give a statistical answer to it. To state our result precisely, we introduce some notation.

Let $n \geq 2$ be fixed, and R be any positive real number. Define $\mathcal{B}_n(R)$ to be the set of bi-Perron algebraic units no larger than R whose characteristic polynomial has degree at most $2n$. Here, by the characteristic polynomial of a bi-Perron algebraic unit λ, we mean the monic palindromic integral polynomial whose leading root is λ and has the lowest degree among all such polynomials. And let $\mathcal{D}_n(R)$ be the set of dilatations no larger than R of pseudo-Anosov maps with orientable invariant foliations on a closed orientable surface S_n of genus n. Similarly, let $\mathcal{D'}_n(R)$ be the set of dilatations no larger than R of pseudo-Anosov maps, not necessarily with orientable invariant foliation, on a closed orientable surface with genus n.

We remark that $\mathcal{D}_n(R)$ is contained in $\mathcal{B}_n(R)$, and $\mathcal{D'}_n(R)$ is contained in $\mathcal{B}_{3n-3}(R)$. A pseudo-Anosov dilatation λ on a surface of genus n is a root of an integral palindromic polynomial of degree at most $2n$ if its invariant foliations are orientable. This is because λ is the leading eigenvalue of the induced symplectic action on the homology group of the surface, which is $\mathbb{Z}^{2n}$. If we do not require the invariant foliation to be orientable, the upper bound on degree is $6n-6$: we can reduce this case to the case of orientable foliation by taking a double cover of the surface, and this bound follows from the fact that a quadratic differential on a surface of genus n has at most $2n$ zeros which is due to Gauss-Bonnet together with the Riemann-Hurwitz formula.

Note that Fried's conjecture is equivalent to $\mathcal{B}_n(R)$ being contained in $\mathcal{D}_m(R)$ for some large enough m. But a priori, m could be arbitrarily large and we do not know how to prove or disprove the claim. Instead we show the following.

Theorem 1. *Let $\mathcal{B}_n(R)$, $\mathcal{D'}_n(R)$ and $\mathcal{D}_n(R)$ be as above. Then*

(1)
$$\lim_{R\to\infty} \frac{|\mathcal{D}_m(R) \cap \mathcal{B}_n(R)|}{|\mathcal{B}_n(R)|} = 0 \ ,$$
where $4m-3 \leq n(n+1)/2$. In particular, $\lim_{R\to\infty} \frac{|\mathcal{D}_n(R)|}{|\mathcal{B}_n(R)|} = 0, \forall n \geq 6$.

(2)
$$\lim_{R\to\infty} \frac{|\mathcal{D'}_m(R) \cap \mathcal{B}_n(R)|}{|\mathcal{B}_n(R)|} = 0 \ ,$$
where $6m-6 \leq n(n+1)/2$. In particular, $\lim_{R\to\infty} \frac{|\mathcal{D'}_n(R) \cap \mathcal{B}_n(R)|}{|\mathcal{B}_n(R)|} = 0, \forall n \geq 10$.

Here $|A|$ means the cardinality of A for a finite set A.

Theorem 1 says that asymptotically almost all bi-Perron algebraic units whose characteristic polynomial has degree at most $2n$ do not correspond to dilatations of pseudo-Anosov maps on a surface of genus n for all $n \geq 10$,

and for $n \geq 6$ if the invariant foliation is further assumed to be orientable. It would be interesting to see if the statement still holds for lower genera.

Let Γ_n be the set of all periodic orbits for the Teichmüller flow on the moduli space of area one abelian differentials on S_n. Then there exists a surjective map $\Gamma_n \to \cup_{R>0}\mathcal{D}_n(R)$ defined by $\gamma \mapsto e^{\ell(\gamma)}$ where $\ell(\gamma)$ is the length of the orbit $\gamma \in \Gamma_n$. Let $\Gamma_n(R)$ be the preimage of $\mathcal{D}_n(R)$ of this map.

By $f \sim g$ we mean $\exists C$ such that $\frac{1}{C}f(x) \leq g(x) \leq Cf(x)$ when $x \gg 0$. By $f \lesssim g$ we mean $f = O(g)$ when $x \to \infty$. At our best knowledge, the following theorem is independently due to Eskin-Mirzakhani-Rafi [5] and Hamenstädt [9] (which is rephrased for our purpose).

Theorem 2 ([5], [9])**.** $|\Gamma_n(R)| \sim \frac{R^{4n-3}}{(4n-3)\log R}$.

We give a brief explanation of the above statement. In [5] and [9], it was stated that when restricted to each connected component of the strata of the moduli space of area one abelian differentials on S_n,

$$\begin{aligned}|\Gamma_n(R)| &= |\{\gamma \in \Gamma_n : e^{\ell(\gamma)} \leq R\}| \\ &= |\{\gamma \in \Gamma_n : \ell(\gamma) \leq \log R\}| \\ &\sim \frac{e^{(2n+\ell-1)\log R}}{(2n+\ell-1)\log R} = \frac{R^{2n+\ell-1}}{(2n+\ell-1)\log R}.\end{aligned}$$

Here ℓ is the maximum number of zeros of an area one abelian differential on S_n, so it is $2n-2$. See [16], [22], [4] and [5] for the relevant background of this theorem.

As a result, we have $|\Gamma_n(R)| \sim \frac{R^{4n-3}}{(4n-3)\log R}$ on each connected component of the strata. But for fixed n, there exists only finite number of such components. Therefore we get $|\Gamma_n(R)| \sim \frac{R^{4n-3}}{(4n-3)\log R}$ without restricting to the components. Note that since n is fixed, we can just say $|\Gamma_n(R)| \sim \frac{R^{4n-3}}{\log R}$.

As a direct corollary, we have:

Corollary 3. $|\mathcal{D}_n(R)| \lesssim \frac{R^{4n-3}}{\log R}$.

In the exactly same way, we can obtain an analogue of Corollary 3 for $|\mathcal{D'}_n(R)|$ from the following theorem which is due to Eskin and Mirzakhani [4]:

Theorem 4 (Theorem 1.1, [4])**.** *The number of geodesics in the moduli space of genus n surface of length at most* $\log(R) \sim \frac{R^{6n-6}}{(6n-6)\log(R)}$.

And as a direct corollary, just like Corollary 3, we have:

Corollary 5. $|\mathcal{D'}_n(R)| \lesssim \frac{R^{6n-6}}{\log R}$.

We remark that this does not imply $|\mathcal{D}_n(R)| \sim \frac{R^{4n-3}}{\log R}$ or $|\mathcal{D'}_n(R)| \sim \frac{R^{6n-6}}{\log R}$, since each element in $\mathcal{D}_n(R)$ or $\mathcal{D'}_n(R)$ may correspond to a lot of different closed geodesics in the moduli space.

Now we study how $|\mathcal{B}_n(R)|$ grows. Let $P_n(R)$ be the set of Perron polynomials of degree n with roots no larger than R, ($x > 1$ is Perron if it is the root of a monic irreducible polynomial with integer coefficients, so that the other roots of the polynomial are (strictly) less than x in absolute value).

Lemma 6. $|P_n(R)| \sim R^{n(n+1)/2}$.

Proof. $|P_n(R)| \lesssim R^{n(n+1)/2}$: Due to Vieta's formula the absolute value of the coefficient of x^k of a monic, degree n polynomial with all roots no larger than R is $\lesssim R^{n-k}$. Hence, the total number of such polynomials must be $\lesssim \prod_k R^{n-k} = R^{n(n+1)/2}$.

$R^{n(n+1)/2} \lesssim |P_n(R)|$: Let a_k be the coefficient of x^k in a degree n monic polynimial (so $a_n = 1$). By Rouché's theorem, when

$$1 > |\frac{a_0}{R^n}| + |\frac{a_1}{R^{n-1}}| + \cdots + |\frac{a_{n-1}}{R}| \tag{1}$$

$$\left(\frac{1}{2}\right)^{n-1} |\frac{a_{n-1}}{R}| > |\frac{a_0}{R^n}| + \cdots + |\frac{a_{n-2}}{R^2}|\left(\frac{1}{2}\right)^{n-2} + \left(\frac{1}{2}\right)^{n} \tag{2}$$

and

$$\left(\frac{1}{3}\right)^{n-1} |\frac{a_{n-1}}{R}| > |\frac{a_0}{R^n}| + \cdots + |\frac{a_{n-2}}{R^2}|\left(\frac{1}{3}\right)^{n-2} + \left(\frac{1}{3}\right)^{n} \tag{3}$$

one root λ of this polynomial has magnitude between $R/2$ and R while all other roots have magnitude smaller than $R/3$. Hence λ must be real. If $\lambda < 0$ one can multiply $(-1)^{n-k}$ to a_k to get a polynomial with a root $-\lambda$. Hence, half of those polynomials have a leading positive real root, so they are in $P_n(R)$.

The inequalities (1), (2) and (3) are satisfied if and only if the point $(a_0/R^n, \ldots, a_{n-1}/R)$ lies in a non-empty open subset $U \subset [-1, 1]^n$. The number of such points as $R \to \infty$ converges to the volume of this open subset divided by the co-volume of the lattice $\mathbb{Z}/R^n \times \mathbb{Z}/R^{n-1} \times \ldots \times \mathbb{Z}/R$, and the co-volume of this lattice is $R^{-n(n+1)/2}$.

□

Lemma 7. $\lim_{R\to\infty} \frac{|\{\textit{reducible elements in } P_n(R)\}|}{|P_n(R)|} = 0$

Proof. Because any reducible monic integer polynomial can be written as the product of two monic integer polynomials of lower degree, we have:

$$|\{\text{reducible elements in } P_n(R)\}| \leq \sum_k |b_k(R)||b_{n-k}(R)|$$

where $b_k(R)$ is the set of monic polynomials with roots bounded by R. The first part of the proof of the previous lemma implies that $|b_k(R)| \lesssim R^{k(k+1)/2}$, hence $|\{\text{reducible elements in } P_n(R)\}| \sim o(R^{n(n+1)/2})$. □

Let $\mathcal{P}_n(R)$ be the set of Perron numbers of degree n no larger than R. They have one-one correspondence with irreducible elements in $P_n(R)$ which by Lemma 7 constitute almost all of $P_n(R)$ asymptotically. Hence by Lemma 6, $|\mathcal{P}_n(R)| \sim R^{n(n+1)/2}$.

Lemma 8. $|\mathcal{B}_n(R)| \sim R^{n(n+1)/2}$.

Proof. When $x > 1$, $1/x < 1$. Hence, $x \in \mathcal{B}_n(R)$ implies that $x + 1/x \in \mathcal{P}_n(R+1)$, hence $|\mathcal{B}_n(R)| \lesssim (R+1)^{n(n+1)/2} \sim R^{n(n+1)/2}$.

On the other hand, from the proof of Lemma 6 and the discussion above, the number of Perron numbers of degree n which lie between R and $R/2$ and have all conjugates smaller than $R/3$ is $\sim R^{n(n+1)/2}$. When R is sufficiently large, each such Perron number y correspond to a bi-Perron number x by the relation $x + 1/x = y$. And in fact x is an algebraic unit (so it is in $\mathcal{B}_n(R)$). Note that both roots of the polynomial $x^2 - yx + 1$ are algebraic integers, since it is a monic polynomial with algebraic integer coefficients. Furthermore, the product of these roots is 1, so they must be algebraic units. Hence $\mathcal{B}_n(R) \gtrsim R^{n(n+1)/2}$. □

Now we are ready to prove our main theorem.

Proof of Theorem 1. By Corollary 3 and Lemma 8, we get

$$\frac{|\mathcal{D}_m(R) \cap \mathcal{B}_n(R)|}{|\mathcal{B}_n(R)|} \lesssim \frac{R^{4m-3}}{\log R \cdot R^{n(n+1)/2}}.$$

The right-hand side goes to 0 as R goes to $+\infty$ as long as we have $4m - 3 \leq n(n+1)/2$. When $m = n$, this inequality is satisfied if and only if $n \geq 6$ (recall that n is always assumed to be at least 2), which proved part (1). Part (2) follows from the same argument above but using Corollary 5 instead of Corollary 3. □

Recall that Γ_n is the set of all closed geodesics in the moduli space of area one abelian differentials on the surface S_n, and $\Gamma_n(R)$ is the subset of Γ_n which consists of the closed geodesics of length no larger than $\log R$. For each $\gamma \in \Gamma_n$, let m_γ be the number $|\{\gamma' \in \Gamma_n : \ell(\gamma') = \ell(\gamma)\}|$.

We remark that if m_γ were uniformly bounded, one could have obtained Theorem 1 (1) for $n \geq 2$ instead of $n \geq 6$ using Theorem 1 (1) of [10]. But at least in the low genus cases, this is not true. As an application of our argument, we obtain the following theorem.

Theorem 9. *Suppose $n \leq 5$. For any positive integer k, the set $\{\gamma \in \Gamma_n : m_\gamma \geq k\}$ is typical, i.e.,*

$$\lim_{R\to\infty} \frac{|\{\gamma \in \Gamma_n(R) : m_\gamma \geq k\}|}{|\Gamma_n(R)|} \to 1.$$

Proof. Suppose not. Then for some k, we have

$$\limsup \frac{|\{\gamma \in \Gamma_n(R) : m_\gamma < k\}|}{|\Gamma_n(R)|} > 0.$$

But this implies that

$$\limsup \frac{|\mathcal{D}_n(R)|}{|\Gamma_n(R)|} \geq \limsup \frac{\frac{1}{k}|\{\gamma \in \Gamma_n(R) : m_\gamma < k\}|}{|\Gamma_n(R)|} > 0.$$

On the other hand, we know that $\mathcal{D}_n(R) \subset \mathcal{B}_n(R)$. As a consequence, $\lim \frac{|\mathcal{B}_n(R)|}{|\Gamma_n(R)|} \geq \limsup \frac{|\mathcal{D}_n(R)|}{|\Gamma_n(R)|}$. By Corollary 3 and Lemma 8, we get

$$\frac{|\mathcal{B}_n(R)|}{|\Gamma_n(R)|} \sim \frac{R^{n(n+1)/2} \log R}{R^{4n-3}} \to 0, \text{ for } n \leq 5,$$

a contradiction.

□

1. Acknowledgements

We would like to thank Balázs Strenner for many helpful comments. The first author was supported by the ERC Grant Nb. 10160104.

References

[1] Arnoux, P., Yoccoz, J.(1980). *Construction de difféomorphisms pseudo-Anosov*, C. R. Acad. Sci. Paris 292, 75–78.

[2] Baik, H., Rafiqi, A. and Wu, C. (2016). *Constructing pseudo-Anosov maps with given dilatations*, Geom. Dedicata, Volume 180, Issue 1, 39–48.

[3] de Carvalho, A. and Hall, T. (2004). *Unimodal generalized pseudo-Anosov maps.* Geom. Topol., 8, 1127–1188.

[4] Eskin, A., Mirzakhani, M. (2011). *Counting closed geodesics in moduli space*, J. Mod. Dynamics 5, 71–105.

[5] Eskin, A., Mirzakhani, M., Rafi, K. *Counting closed geodesics in strata*, arXiv:1206.5574.

[6] Farb, B., Margalit, D. (2012). *A primer on mapping class groups*, volume 49 of Princeton Mathematical Series. Princeton University Press, Princeton, NJ.

[7] Fried, D. (1985). *Growth rate of surface homeomorphisms and flow equivalence.* Ergod. Theory Dyn. Syst., 5(04), 539–563.

[8] Hamenstädt, U. *Symbolic dynamics for the Teichmüller flow*, arXiv:1112.6107.

[9] Hamenstädt, U. (2013). *Bowen's construction for the Teichmüller flow*, J. Mod. Dynamics 7, 489–526.

[10] Hamenstädt, U. *Typical properties of periodic Teichmüller geodesics*, arXiv:1409.5978.

[11] Hironaka, E. (2010). *Small dilatation mapping classes coming from the simplest hyperbolic braid*, Algebr. Geom. Topol., 10(4):2041–2060.

[12] Lanneau, E., Thiffeault, J. (2011). *On the minimum dilatation of braids on punctured discs*, Geom. Dedicata, 152:165–182.

[13] Lanneau, E., Thiffeault, J. (2011). *On the minimum dilatation of pseudoAnosov homeromorphisms on surfaces of small genus*, Ann. Inst. Fourier (Grenoble), 61(1):105–144.

[14] Leininger, C. (2004) *On groups generated by two positive multi-twists: Teichmuller curves and Lehmer's number*, Geom. Topol., 8:1301–1359.

[15] Long, D. (1985). *Constructing pseudo-Anosov maps*, In Knot theory and manifolds (Vancouver, B.C., 1983), volume 1144 of Lecture Notes in Math., 108–114. Springer, Berlin.

[16] Masur, H. (1982). *Interval exchange transformations and measured foliations*, Ann. Math. 115, 169–201.

[17] Penner, R. (1988). *A construction of pseudo-Anosov homeomorphisms*, Trans. Amer.Math. Soc., 310(1):179–197.

[18] Shin, H., Strenner, B. (2015). *Pseudo-Anosov mapping classes not arising from Penner's construction*, Geom. Topol. 19, 3645–3656.

[19] Thurston, W. (1988). *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Am. Math. Soc. (New Ser.) 19, 417–431.

[20] Thurston, W. (2014). *Entropy in dimension one.* arXiv:1402.2008.

[21] *Travaux de Thurston sur les surfaces*, volume 66 of Asterisqué . Société Mathématique de France, Paris, 1979. Séminaire Orsay, With an English summary.

[22] Veech, W. (1986). *The Teichmüller geodesic flow*, Ann. Math. 124, 441–530.

Mathematisches Institut, Rheinische Friedrich-Wilhelms-Universität Bonn, Endenicher Allee 60, 53115 Bonn, Germany
E-mail address: baik@math.uni-bonn.de

Department of Mathematics, Cornell University, Malott Hall, Ithaca, NY 14853, USA
E-mail address: ar776@cornell.edu

Max Planck Institute for Mathematics, Bonn, Germany
E-mail address: chenxiwu@mpim-bonn.mpg.de